



## Original Research Article

## Hypoglycemic and antidiabetic evaluation of leaves of phyllanthus distichus in normal and alloxan induced diabetic rats

Debashis Mohapatra<sup>1,\*</sup>, Sunita Mishra<sup>1</sup>

<sup>1</sup>College of Pharmaceutical Sciences, Baliguali, Puri, Odisha, India



## ARTICLE INFO

## Article history:

Received 18-08-2020

Accepted 22-08-2020

Available online 23-10-2020

## Keywords:

Alcoholic Extract of Phyllanthus

Distichus (AEPD)

Antigen (Ag)

Alanine Amino Transferase (ALAT)

Alkaline Phosphate (ALP)

Aspartate Aminotransferase (ASAT)

Aqueous (Aq)

Body Wt

Direct Bilirubin (DB) Diabetes

Mellitus (DM)

Phyllanthus Distichus (PD)

## ABSTRACT

The ethanolic extract of leaves of Phyllanthus distichus results maximum yield value than that of petroleum ether extract, chloroform extract through successive extraction process. The ethanolic extract of leaves of Phyllanthus distichus showed maximum control in blood sugar in hyperglycemic wistar rats than other experimental extracts. The test extract also reduces the blood sugar level to a maximum extent in case of normal animals. Among the study of effects of alcoholic extracts of leaves of Phyllanthus distichus in both normoglycemic and hyperglycemic model through oral route. Toxicological study revealed that the alcoholic extracts Phyllanthus distichus leaves were safe and does not alter normal physiological and behavioral effect even at a higher dose level of 3000 mg/kg body weight.

Administration of alcoholic extracts Phyllanthus distichus significantly reduces the elevated glucose level in alloxan induced diabetic rats confirms its anti-diabetic activity. This also reduces normal glucose level, which reveals the hypoglycemic property. The whole protein, whole cholesterol, ASAT, ALAT, ALP enzyme activity of alloxan administered rats showed significantly higher than normal rats, the test extract treated rats significantly reduced the elevated levels. The results of the present investigation indicate that the alcoholic extracts Phyllanthus Distichus leaves may have a place in the therapy of Diabetes Mellitus as anti-diabetic and/or hypoglycemic agent. The hypoglycemic and/or anti-diabetic effect of the test extract may be due to the influence on glycogenesis, glycogenolysis metabolic activity property of one or more of its constituents. Ethanolic extract of leaves of Phyllanthus Distichus is beneficial in lowering the blood sugar concentration and in the management of other diabetic complications without any doubt.

© 2020 Published by Innovative Publication. This is an open access article under the CC BY-NC license (<https://creativecommons.org/licenses/by-nc/4.0/>)

## 1. Introduction

The studies on the leaves of P.D. have not been reported on the antidiabetic activity. As for the above reason the present study was carried out on mice and rats and the dose fixation was done, as per the protocol.

1. Collection of leaves of P.D. and extraction of processes using different solvents according to their increasing order of polarity, for the present investigation.
2. Determination of the most potent extract for reduction of the elevated blood glucose levels induced by Alloxan in wistar rats by single dose treatment model through measuring the fasting blood sugar levels.

3. Acute toxicity study of the potent extract of P.D to determine LD<sub>50</sub> and selection of dose.
4. The route of administration in oral route by single dose treatment in both normoglycemic and hyperglycemic model.
5. Diabetes is induced by rat by single dose Alloxan administration.
6. Diabetes is confirmed by blood glucose estimation.
7. The study of hypoglycemic and/or antidiabetic activity of the most potent extract is to be done by the following models.

## 1.1. Effect of test extracts on

1. Single dose treated normoglycemic rats.
2. Single dose treated hyperglycemic rats.

\* Corresponding author.

E-mail address: [dmohapatra911@gmail.com](mailto:dmohapatra911@gmail.com) (D. Mohapatra).

3. Single dose treated glucose loaded hyperglycemic rats.
4. Multi dose treated normoglycemic rats.
5. Multi dose treated hyperglycemic rats.
6. To estimate the glycogen conc. In liver and kidney.
7. Behavioural changes of animals after single dose treatment if any also to be performed.
8. Plasma lipids(TC,TG, LDL,HDL,VLDL) are to be analysed in diabetic rats to find out wheather any correction exits between the levels of these parameters and diabetes.
9. The changes in the levels of alkaline phosphate, aminotransferase(ASAT,ALAT), Total proteins, total bilirubin, direct bilirubin, albumin and globulin etc. are to be analyse in a serum.
10. Haematological parameters to be studied in the last day of treatment of subacute toxicity study.
11. Determination of percentage loss of body weights in alloxan induced diabetic rats treated with test extract at the end of 21 days.

The results obtained from control and experimental animals of the present investigation are compared and discussed in the light of available literatures.

## 2. Materials and Methods

### 2.1. Plant collection and authentication

Large number of leaves of *Phyllanthus distichus* appears in the month of July and November. Leaves were collected from the interior villages of Athgarh (Orissa). Before their use, they were carefully identified and collected.

### 2.2. Preparation of the powder

The leaves were dried over a polythene cover in shade drying method with a help of fan at 21 °C room temperature and pulverized using a mechanical grinder. The powder is further passed through a fine mesh sieve to get a fine powder.

### 2.3. Preparation of the extracts

The powdered material (600gm) was extracted separately with different solvents of increasing order of polarity, e.g.. Petroleum ether, chloroform and ethanol by soxhletion method. The liquid extracts were concentrated under vaccum to yield dry extracts and preserved in a desicator till further experiments.

### 2.4. Percentage yield of the extracts

The leaves of *Phyllanthus distichus* was extracted with solvents of increasing polarity by Soxhlet apparatus. The percentage yield of the leaves of *Phyllanthus distichus* was found to be 2.66%, 1.66%, 14.66% and 12.66% with petroleum ether, chloroform, alcohol and water, respectively(shown in tab-1). The percentage yield of the

ethanolic extract of *Phyllanthus distichus* was found to be greater (14.66%) than extract with other solvent.

### 2.5. Preparation of drug solution

#### 2.5.1. Preparation of extract solution

The semisolid extracts of P.D. and dissolved in sufficient quantity of solvent (Tween & Normal saline). The aqueous extract of P.D. was dissolved in Normal saline and the latter were used for administration.

#### 2.5.2. Glibenclamide

Glibenclamide 2.5 mg/kg b.wt was dissolved in sufficient quantity of solvent (Normal saline) and used in the treatment.<sup>1</sup>

#### 2.5.3. Alloxan

Alloxan 120mg/kg b. wt was dissolved in the Normal saline and used to induce diabetes in rats.<sup>2</sup>

### 2.6. Selection of aAnimals

Healthy adult albino rats of wister strain weighing 150-200 gms and adult albino mice of swiss strain weighing 20-25 gms were selected for toxicological and anti-diabetic studies respectively with the approval from the instutional animal ethics committee (Regd. No.1171/C/08/CPCSEA).

### 2.7. Maintenance of animals

The animal house was well ventilated and animals had 12± 1hour day and night schedule with temperature between 11-20±2°C. The animals were housed in large spacious hygienic cages during the course of the experimental period. The animals were fed with rat pellets feed supplied by M/s Hindustan Lever Ltd., Bangalore, India and Aquaguard filtered water ad libitum. The place where the experiments we conducted kept very hygienic by cleaning with antiseptic solutions. The husk, which was for the purpose of keeping as a bed to the animals, was autoclaved and cleaned. Before the animals were kept, the cages (polypropylene) were also sterilized along with the water feeding bottles. As the diabetic animals are susceptible to infection hygienic condition were necessary.

## 3. Experimental Setup

### 3.1. Hypoglycemic activity of AEPD in single dose treatment

#### 3.1.1. In normoglycemic animal

To check wheather the alcoholic extract of P.*distichus* is acting as an insulin secretagogue (Stimulation of insulin secretion from the pancreatic  $\beta$ -cell & to check the utilization). The study was carried out as mentioned Table 1.

**Table 1:** Overnight fasted animals were grouped in 4 groups of 6 animals each groups

G roup	Animal used	S ex	Body wt.	T reatment
Gr-I	Albino rat	M	200gms	Served as control(Normal saline). Once daily
Gr-II	Albino rats	M	200gms	Served as standard(Glibenclamide 2.5mg/kg)in p.o. Once daily
Gr-III	Albino rats	M	200gms	Test drug(p.o/i.p) Dose -1. Once daily
Gr-IV	Albino rats	M	200gms	Test drug(p.o/i.p) Dose-2. Once daily

Blood glucose levels were measured in all 5 groups in diff. Time interval like 1, 2, 4, 8, 12hrs.

### 3.1.2. Alloxan induced diabetic animal

To check wheather the Ethanolic extract of *P.distichus* has antidiabetic activity. The study was carried out mentionedTable 2.

### 3.1.3. Glucose loaded hyperglycemic animals

It is a standard procedure used to monitor blood glucose levels over a period of time following glucose load.

To check wheather the ethanolic extract of *P.distichus* has hypoglycemic activity as mentioned Table 3.

## 3.2. Anti diabetic & hypoglycemic study of extract of *p.distichus* in multi dose treatment

### 3.2.1. In normoglycemic animals

To check wheather the ethanolic extract of *P.distichus* has hypoglycemic activity on the multiple dose treatment.

### 3.2.2. In diabetic animals

To check wheather the extract of *P.distichus* has antidiabetic activity on the multiple dose treatment in alloxan induced animals; the study was carried as mentioned Table 5.

## 3.3. Toxicity studies

### 3.3.1. Acute oral toxicity studies

The acute toxicity study is aimed to establish the therapeutic index i.e., the ratio between the pharmacologically effective dose and the lethal dose, and also to perform the primary screening.

The aqueous extract of *Phyllanthus distichus* was administered once orally at 4 dose levels (500, 1000, 2000,3000 mg/kg) to group of 10 mice of both sexes about equal in number which have been fasting overnight (about 18 hrs.)

The treated mice were observed continuously for two hours and then occasionally for further for four hours and finally overnight mortality recorded.<sup>3</sup>

During the course of study the behaviours of the mice were carefully observed and fall of time, reduction of spontaneous activity also determined.<sup>4</sup>

### 3.3.2. Sub acute toxicity studies

The purpose of this study is to determine the maximum tolerated dose and daily dosed for 3 to 4 weeks and to indicate the nature of toxic reaction of the drug.

In present study, the influence of *Phyllanthus distichus* extract on haematological parameters, a pathological change on 21 days dosing in Alloxan induce diabetic rats was carried out.

The diabetic rats were grouped into 6 of 6 animals each. Group 1 animals received solvent, Group III and Group IV was received AEPD and 200 mg/kg b. wt. p.o., once daily for 21 days.

At the end of the experimental period the blood was collected, serum separated and subjected to hematological and biochemical examination.<sup>3</sup>

## 3.4. Investigation carried out in the present study

Blood sample was drawn into tubes at different time intervals.

Serum was separated out by centrifugation and used in different estimations.

Diabetes was induced by a single dose of alloxan administration.

Blood glucose level, which is elevated in association with hyperglycemia diabetes, was measured to assess whether the plant extract has antidiabetic/hypoglycemic effect.

A significant body weight reduction is associated with diabetes hence the body weight was recorded before, during and after the treatment.

Acute toxic study was carried out to find out toxic symptoms of the extract and gross behavioural changes were also noticed.

Sub acute toxicity study was carried out to find out the influence of the extract on biochemical, haematological and histopathological findings.

ALAT, ASAT, ALP etc. . . were assayed in serum.

Serum lipids like total Cholesterol, HDL, LDL and VLDL, triglyceride were estimated.

Serum total protein and albumin were analysed.

Blood parameters like haemoglobin, RBC, WBC etc. . . . counts were determined

Clotting time was also determined

**Table 2:** Overnight fasted animals were grouped in 4 groups of 6 animals each groups.

Group	Animal used	Sex	Body wt.	T treatment
Gr-I	Albino rat	M	200gms	Served as control(Normal saline). Once daily
Gr-II	Albino rats	M	200gms	Served as standard(Glibenclamide 2.5mg/kg)in p.o. Once daily
Gr-III	Albino rats	M	200gms	Test drug(p.o/i.p) Dose -1. Once daily
Gr-IV	Albino rats	M	200gms	Test drug(p.o/i.p) Dose-2. Once daily

Blood glucose levels were measured in all 5 groups in diff. Time interval like 1, 2, 4, 8, 12hrs.  
(All these Rats are treated with Alloxan 110-150mg/kg i.p.prior to 72 hrs)

**Table 3:** Overnight fasted animals were grouped in 4 groups of 6 animal each groups & glucose (1gm/kg) was loaded to group II & IV

G roup	Animal used	Sex	Body wt.	T treatment
Gr-I	Albino rat	M	200gms	Served as control(Normal saline). Once daily
Gr-II	Albino rats	M	200gms	Served as standard(Glibenclamide 2.5mg/kg)in p.o. Once daily
Gr-III	Albino rats	M	200gms	Test drug(p.o/i.p) Dose -1. Once daily
Gr-IV	Albino rats	M	200gms	Test drug(p.o/i.p) Dose-2. Once daily

Blood glucose levels were measured in all 5 groups in diff. Time interval like 1, 2, 4, hrs.

**Table 4:** Overnight fasted animals were grouped in 4 groups of 6 animal each groups

G roup	Animal used	Sex	Body wt.	Treatment
Gr-I	Albino rat	M	200gms	Served as control(Normal saline). Once daily
Gr-II	Albino rats	M	200gms	Served as standard(Glibenclamide 2.5mg/kg)in p.o. Once daily
Gr-III	Albino rats	M	200gms	Test drug(p.o/i.p) Dose -1. Once daily
Gr-IV	Albino rats	M	200gms	Test drug(p.o/i.p) Dose-2. Once daily

Blood glucose levels & body wt. were measured on 0<sup>th</sup>, 3<sup>rd</sup>, 9<sup>th</sup> day

**Table 5:** Overnight fasted animals were grouped in 4 groups of 6 animal each group

Group	Animal used	Sex	Body wt.	T treatment
Gr-I	Albino rat	M	200gms	Served as control(Normal saline). Once daily
Gr-II	Albino rats	M	200gms	Served as standard(Glibenclamide 2.5mg/kg)in p.o. Once daily
Gr-III	Albino rats	M	200gms	Test drug(p.o/i.p) Dose -1. Once daily
Gr-IV	Albino rats	M	200gms	Test drug(p.o/i.p) Dose-2. Once daily

Blood glucose levels & body wt. were measured on 0<sup>th</sup>, 7<sup>th</sup>, 14<sup>th</sup> & 21<sup>st</sup> day.

### 3.5. Induction of diabetes

In the present study a single dose of Alloxan in normal saline 120 mg/kg b. wt was administered intraperitoneally.<sup>2</sup>

Diabetes developed gradually was assessed after a week an experiment was carried out to determine the blood sugar levels. Animals with blood sugar levels 200-250 mg/dl, were chosen on 7<sup>th</sup> day and considered as diabetic rats for antidiabetic screening.

## 4. Blood Collection

A small amount of blood without sacrificing the animal by orbital sinus puncture. The rats were made semi conscious with ether using the sterile capillary tube, puncture the

orbital sinus at the inner canthus of the eye, by rotating the capillary tube with sufficient but not excessive pressure, two or three times.

As bleeding starts, the animal was held close to the centrifuge tube, and blood was collected. Applying pressure on the inner canthus for a short while stopped the bleeding. The collected blood was used to determine the biological parameters.<sup>5</sup>

### 4.1. Determination blood glucose levels

A small amount of blood without sacrificing the animals by orbital sinus puncture or by snipping off the tip of the tail. The rats were made semi-conscious with ether using the

sterile blunt needle; puncture the orbital sinus at the inner canthus of the eye, by rotating the needle with sufficient but not excessive pressure, two or three times as described in sub-acute toxicity study.

As bleeding starts, the animal was held close to the haemogluco test strip and allows the drop of blood to fall on the strip. The bleeding was stopped by applying pressure on the inner canthus for a short while.<sup>5</sup>

The Johnson & Johnson Ltd. (One touch – Life Scan) glucometer was switched on. When the instrument gives a beep sound after 1 min. The test strip was inserted. Then as bleeding starts, the animal was held close to the haemogluco test strip and allows the drop of blood to fall on the strip. Then the instrument was allowed to react for one minute. Then the blood glucose level was displayed on the screen was recorded.

#### 4.2. Body weight analysis

Initial body weight of rats was recorded. The final body weight of rats was recorded on 3<sup>rd</sup> day, 7<sup>th</sup> day, 14<sup>th</sup> day, and 21<sup>st</sup> day. The % loss in body weight was calculated after 21 days and the weight variation was noted from treated groups and compared with normal, Alloxan control group.

#### 4.3. Preparation of serum

The collected blood was kept at room temperature, refrigerated and centrifuged, for 20 min at 2000 rpm to separate serum.

#### 4.4. Biochemical parameters estimated

Asparate aminotransferase (ASAT)  
 Alanine Aminotransferase (ALAT)  
 Alkaline Phosphatase (ALP)  
 Total Cholesterol (Tc)  
 High Density Lipid Cholesterol (HDL-C)  
 Triglyceride (TG)  
 Total Protein (Tp)  
 Albumin  
 Globulin  
 Total Bilirubin (TB)  
 Direct Bilirubin (DB)

#### 4.5. Transaminases

##### 4.5.1. Asparate aminotransferase(ASAT):<sup>6,7</sup>

Asparate aminotransferase in serum was assayed using Ecoline dignostic kit. ASAT analyses the following reactions. Glutamate + Oxalacetate Malate + NAD.

The rate of NADH consumption was measured photometrically at 340nm and is directly proportional to the ASAT activity in the sample.

Asparate aminotransferase level in serum in expressed as U/L. Glutamine + Pyruvate Lactate + NAD.

##### 4.5.2. Alanine aminotransferase in serum was assayed using Ecoline dignostic kit. ALAT analyses the following reaction

The rate of NADH consumption was measured photometrically at 340nm and is directly proportional to the ALAT activity in the sample.

Asparate aminotransferase level in serum in expressed as U/L.

#### 4.6. Alkaline phosphatase (ALP)

Alkaline Phosphatase in serum was assayed using Ecoline dignostic kit. ALP analyses the following reaction. Phosphate + 4-nitrophenolate

The rate of increase of 4-nitrophenolate was determined photometrically at 405 nm and is directly proportional to the ALP activity in the sample.

#### 4.7. Total cholesterol (Tc):<sup>8,9</sup>

Cholesterol in serum was estimated by CHOD-PAP method using an Ecoline Diagnostic Kit. Cholesterol and its esters are released from lipoproteins by detergents. Cholesterol esterase hydrolyses the esters. In the subsequent enzymatic oxidation by cholesterol osidase, H<sub>2</sub>O<sub>2</sub> is formed. This converted into a coloured quinonimine in a reaction with 4 aminoantipyrine and phenol catalyzed by peroxidase. The absorbance of the sample and of the standard was measured against the reagent blank value at 546mm.

Cholesterol level in serum/liver and aorta was expressed as mg/dl or mg/g tissue.

#### 4.8. HDL cholesterol

The HDL cholesterol was separated from Serum after precipitation of LDL and VLDL cholesterol by phosphotungstic acid precipitating reagent.

The supernatant after centrifugation was estimated using Ecoline diagonostic kit by CHOD-PAP method.<sup>8,9</sup>

The absorbance of the sample of the standard was measured against the reagent blank value at 546 mm.

HDL Cholesterol level in serum was expressed as mg/dl.

#### 4.9. VLDL cholesterol

Cholesterol was calculated by the formula<sup>10</sup>

$$\text{Cholesterol} = \text{Triglycerides}/5$$

Cholesterol level in serum was expressed as mg/dl

#### 5. LDL cholesterol

Cholesterol was calculated by using the formula<sup>10</sup>

$$\text{Cholesterol} = \text{Total Cholesterol} - (\text{HDL Cholesterol} - \text{Triglycerdes}/5)$$

Cholesterol level in serum was expressed as mg/dl

### 5.1. Triglycerides (TG):<sup>11</sup>

Triglyceride level in serum/liver and aorta was estimated using an Ecoline Diagnostic kit. Glycerol + Fatty acid Glycerol – 3 phosphate + HO

The absorbance of the sample and of the standard was measured against the reagent blank value at 546 nm. Triglyceride level in serum/liver and aorta was expressed as mg/dl or mg/g tissue.<sup>8,11</sup>

Total protein in serum was assayed using Ecoline diagnostic kit. Proteins and peptides in contrast to other nitrogen containing compounds (e.g. creatinine, urea, uric acid.) produce a violet colored complex with copper ions in an alkaline solution. The so-called burette reaction is particularly easy to carry out, giving reproducible results, which are in good agreement with Kjeldahl method.

Measured the absorbance of the sample of the standard against the burret reagent and the absorbance of the blank against water of 546 nm. Total protein level in serum is expressed as mg/dl.

### 5.2. Albumin<sup>8,12</sup>

Albumin in serum was assayed using Ecoline diagnostic kit.

Albumin forms Blue-green complex with bromocresol green at slightly acidic PH which is measured photometrically the absorbance of the sample and of the standard against blank at 540-600 nm.

Albumin level in serum is expressed as g/dl is recorded.

### 5.3. Globulin

Globulin concentration in serum is indirectly determined by subtracting the albumin concentration from the total protein concentration.<sup>13</sup>

### 5.4. Total bilirubin

Total bilirubin level in serum was assayed using an Ecoline Diagnostic Kit. The total bilirubin in serum is determined by coupling with diazotized sulfanilic acid after the addition of caffeine. Sodium benzoate and sodium acetate. A blue azobilirubin is formed in alkaline Fehlings solution II. This blue compound can also be determined selectively in the presence of yellow by-products (Green mixed coloration) by photometry at 578 nm.

### 5.5. Direct bilirubin<sup>14</sup>

Direct bilirubin in serum was assayed using Ecoline diagnostic kit. The direct bilirubin is measured as the red azo dye at 546 nm without the addition of alkali. This method is based on the definition of direct bilirubin as the quantity of bilirubin. Which, without the addition of an accelerator, can be determined after a reaction time of 5 minutes? This bilirubin comprises mainly the water soluble bilirubin glucuronides. Under the condition used here, free

bilirubin reacts slowly.

Measured the absorbance of sample against blank at 546 nm.

### 5.6. Estimation of liver glycogen

The liver exercised from sub acute toxicity study was subjected to estimate the glycogen.

### 5.7. Procedure

Known weight of the liver tissue was subjected to alkali digestion with 30% KOH in boiling water bath for 20min. 3.0ml of ethanol was added and tubes were kept in a freezer overnight. They were centrifuged at 3000 rpm for 40 min. The precipitate was dissolved in warm water, reprecipitated with ethanol and centrifuged again. The final precipitate was dissolved in 3.0 ml of distilled water and heated for 5 min. in a boiling water bath. Aliquots of the sample were mixed with 4.0ml of Anthrone reagent, heated in a boiling water bath for 20min. The green colour developed was read at 600nm using systronics UV-VIS spectrophotometers.<sup>15</sup> The glycogen content in the tissue is expressed as mg/g wet tissue.

#### 5.7.1. Determination of Hematological parameters

Hemoglobin concentration in the blood sample was estimated as per the standard procedure.<sup>16</sup> The Hb concentration in blood was expressed as gm %.

#### 5.7.2. Total W.B.C. count

Estimation of W.B.C. count was carried out as per the standard procedures.<sup>16</sup> The W.B.C. count was expressed as number of cells/ $\mu$ l of whole blood.

#### 5.7.3. Total R.B.C. count

Estimation of R.B.C. count was carried out as per the standard procedures (Ghai 1988).<sup>17</sup> The R.B.C. count was expressed as number of cells/ $\mu$ l of whole blood.

### 5.8. Clotting time

Clotting time was determined as per the standard procedure by Ghai (1988).<sup>15</sup> It was expressed in minutes.

### 5.9. Statistical analysis

The values are represented as  $\pm$  S.E.M.

The data obtained from various studies were subjected to one-way analysis of variance.(ANOVA)<sup>18</sup> followed by Dunnet's t-test

## 6. Experimental Results & Discussion

1. The present study was divided into 9 phases.

2. In the first phase the *Phyllanthus distichus* leaves are collected, shed dried and the extracts are prepared using different solvents of increasing polarity by the method of Soxhlet extraction.
3. In the second phase the most potent extract for reduction of blood sugar level in alloxan induced hyperglycemic rats was determined.
4. In the third phase the acute toxicity study was done to determine LD50,
5. In the fourth phase the route of administration was determined in single dose normo/hyperglycemic model.
6. In the fifth phase the hypoglycemic effect and anti diabetic effect of the most potent extract was carried out in normoglycemic, glucose loaded hyperglycemic and alloxan induced diabetic rats in single dose treatment.
7. In the sixth phase the hypoglycemic effect and anti diabetic effect of the extract was carried out in normoglycemic and alloxan induced diabetic rats in multi dose treatment or daily dosing for 21 days study.
8. In the seventh phase the percentage loss in the body weight was calculated and analysed after 21 days in the sub acute toxicity study.
9. In the eighth phase the effect of the extract on the glycogen concentration in liver and kidney was estimated.
10. In the ninth phase the effect of the extract in long term use if any, was carried out by observing the various biochemical parameters, haematological profiles, lipid profiles.

#### 6.1. Table I Effect of various extracts of leaves of PD on Alloxan induced diabetic rats

The table I shows that the chloroform extract, aqueous extract & ethanolic extract registered 3.63%, 15.51%, 49.84% decrease in fasting blood glucose levels respectively at the dose level of 100mg/kg body wt. in oral route in alloxan induced diabetic rats at the end of 8hrs. However at the same time solvent control (distilled water) & petroleum ether showed gradual increase in blood sugar level. The standard drug Glibenclamide (2.5mg/kg bwt) possesses 60.97% decrease in blood glucose level at the same time.

The aqueous extract significantly decreases the blood sugar level at the end of 4hrs & 8hrs in the level of  $P < 0.01$ , while the ethanolic extract showed significant decrease of blood sugar level starting from 2hrs to the end of 8hrs at the level of  $P < 0.05$  to  $P < 0.001$  when compared with the solvent control. However the standard drug decrease a significant decrease of blood sugar level beginning from 1hr ( $P < 0.05$ ) up to 8hr ( $P < 0.001$ ) when compared with solvent.

The blood glucose lowering property of ethanolic extract of PD is comparable with the standard drug Glibenclamide (2.5mg/kg Body wt).

Among the tested extract the ethanolic extract at the dose level of 100mg/kg through oral route possess highest percentage decrease of blood sugar level.

Therefore the ethanolic extract was selected for the evaluation of hypoglycemic &/or anti-diabetic activity.

#### 6.2. Acute toxicity study of ethanolic extract of leaves of PD

The result shown in the Table 7 indicated that the ethanolic extract of PD showed no change of behaviour up to 2hrs & no motility was observed up to 24 hrs at the maximum dose levels of 3000mg/kg. Therefore further studies were carried out at the dose levels of 100mg/kg, 200mg/kg, & 300mg/kg body wt.

#### 6.3. Hypoglycemic activity of ethanolic extract of leaves of PD in single dose treatment in normoglycemic rats in oral route

The data recorded in the Table 8 showed the test extract possesses 27.11% & 33.8% decrease in blood glucose level at the dose level of 100mg/kg & 200mg/kg b.wt. respectively in oral route at the end of 8hrs. While the standard drug Glibenclamide (2.5mg/kg) registered 46.9% reduction at the same time. However the solvent control group showed negligible reduction of blood glucose level at the end of 8<sup>th</sup> hrs. The ethanolic extract at the dose level of 100mg/kg showed significant decrease ( $P < 0.01$ ) at 8hrs while at the same time the dose level 200mg/kg showed more significant ( $P < 0.001$ ) when compared to solvent control.

#### 6.4. Anti diabetic activity of ethanolic extract of leaves of PD in single dose treatment in alloxan induced Hyperglycemic rats in oral route

The result expressed in the table iv revealed that the Ethanolic extract in the dose level of 100mg/kg & 200mg/kg body wt possess 49.67% & 55.48% reduction in the blood glucose level respectively at the end of 8hr while the standard drug Glibenclamide (2.5mg/kg body wt.) registered 61.01% decrease in blood sugar level at the same time.

The test extract showed a significant decrease ( $P < 0.001$ ) of blood glucose level starting from 4<sup>th</sup> hr onward in the tested dose levels while standard drug possess the same rate of significance beginning from 2hrs while compared with solvent control.

#### 6.5. Hypoglycemic activity of ethanolic extract of leaves of PD in single dose treatment in glucose loaded hyperglycemic rats in oral route. (OGTT)

The data recorded in table-V indicated that the test extract at the dose level of 100mg/kg & 200mg/kg b.wt. registered 19.45% & 23.55% decrease in blood glucose level at the end

**Table 6:** Effects of various Extracts of leaves of P.D on alloxan induced diabetic rats

Group & treatment	Blood glucose levels(mg/dl)					% age decrease at the end of 8hrs
	0hrs	1hrs	2hrs	4hrs	8hrs	
I.Solvent (saline Water)	254.16±10.55	256±12.79	277.16±9.3	289.16±9.3	294±10.25	
II.Standard Glibenclamide(2.5mg/kg)	246.6±13.9	223±11.45*	200.16±15.56***	135.33±8.4***	96.16±6.3***	60.97
III.Pet Ether Extract (100mg/kg)	264±10.3	270±10.07	279.67±9.93	281.5±5.19	278.5±6.62	
IV.Chloroform Extract (100mg/kg)	275.3±4.03	272±6.08	273.16±8.14	266.67±8.1	265.84±7.6	3.63
V. Ethanol Extract (100mg/kg)	274.83±6.65	264.5±6.4	223.34±11.27**	164.83±12.62***	137.84±7.95***	49.84
VI.Aquous Extract (100mg/kg)	277.16±9.7	267.5±7.3	249±6.9	246.34±91**	234.16±4.6**	15.51
VII. F values		2.26 <sup>b</sup>	9.12 <sup>b</sup>	50.93 <sup>b</sup>	119.75 <sup>b</sup>	

Values are expressed in MEAN ± S.E.M of six animals. One Way ANOVA followed by Dunnet's t-test.

(F-value denotes statistical significance at (ap<0.05, bp<0.01)

(t-value denotes statistical significance at \*p<0.05, \*\*p<0.01 and \*\*\*p<0.001 respectively, in comparison to group-I.

**Table 7:** Acute toxicity study of ethanolic extract of leaves P.D

Group	Dose	Route	No. of Animals	Death
I	500	ORAL	4	0
II	1000	ORAL	4	0
III	1500	ORAL	4	0
IV	2000	ORAL	4	0
V	3000	ORAL	4	0

**Table 8:** Hypoglycemic activity of ethanolic extract of leaves of P.D in single dose treatment in normoglycemic rats in oral route.

Group & treatment	Blood glucose levels(mg/dl)					% age decrease at the end of 8hrs
	0hrs	1hrs	2hrs	4hrs	8hrs	
I. Solvent (saline Water)	104.16±2.21	104.83±1.8	102.83±2.08	106±1.59	105.66±1.5	
II. Standard Glibenclamide(2.5mg/kg)	103±2.25	90±3.02**	84.83±2.03***	79±1.5***	54.67±3.67***	46.9
III. Test-1 (100mg/kg)	104.5±2.01	104.5±2.4	97.33±1.89	94.66±1.96**	76.17±5.14***	27.11
IV. Test-2 (200mg/kg)	104.5±1.98	104.16±1.6	96.16±2.27*	87.16±1.9***	68.84±3.47***	33.8
V. F Values		9.94 <sup>b</sup>	13.22 <sup>b</sup>	42.01 <sup>b</sup>	34.009 <sup>b</sup>	

Values are expressed in MEAN ± S.E.M of six animals. One Way ANOVA followed by Dunnet's t-test.

(F-value denotes statistical significance at (ap<0.05, bp<0.01)

(t-value denotes statistical significance at \*p<0.05, \*\*p<0.01 and \*\*\*p<0.001 respectively, in comparison to group-I.



**Table 9:** Antidiabetic activity of ethanolic extract of leaves of P.D in single dose treatment in Alloxan induced Hyperglycemic rats in oral route

Group & treatment	Blood glucose levels(mg/dl)					%age decrease at the end of 8hrs
	0hrs	1hrs	2hrs	4hrs	8hrs	
I. Solvent (saline Water)	254±10.57	256±12.79	277.16±10.97	289.16±9.3	294±10.25	
II. Standard Glibenclamide(2.5mg/kg)	246.67±13.9	234±12.17	216.83±18.22**	135.34±8.4***	96.16±6.3***	61.01
III. Test-1 (100mg/kg)	274.83±6.6	264.16±6.4	235.5±5.23*	168.17±5.61***	138.34±3.09***	49.67
IV. Test-2 (100mg/kg)	273.34±5.4	260.16±5.1	228.33±4.3	145.34±5.4***	121.67±12.2***	55.48
V. F Values		1.9	5.5 <sup>b</sup>	92.06 <sup>b</sup>	104.88 <sup>b</sup>	

Values are expressed in MEAN ± S.E.M of six animals. One Way ANOVA followed by Dunnet's t-test.

(F-value denotes statistical significance at (ap<0.05, bp<0.01)

(t-value denotes statistical significance at \*p<0.05, \*\*p<0.01 and \*\*\*p<0.001 respectively, in comparison to group-I.

of 4hrs respectively. While the standard drug Glibenclamide (2.5mg/kg) reduces blood glucose level up to 47.28% at the same time.

At the end of 4hrs. the test extract at both the dose levels bears a significant (p<0.001) decrease of blood glucose in compared with solvent control. While the standard drug Glibenclamide holds the same significance (p<0.001) from the beginning from 2hrs

It is observed that the AEPD leaves effectively control the blood sugar level in normal as well as glucose loaded hyperglycemic rats in single dose treated model. Such an effect might be decrease in the rate of initial glucose absorption when the plant fiber is given orally with glucose.<sup>18</sup> Hence it may be presumed that the glucose lowering effect of the extract was achieved by an extra intestinal ac

#### 6.6. Hypoglycemic activity of ethanolic extract of leaves of PD in multidose treatment in normoglycemic rat in oral dose

The purpose of the study is to establish the therapeutic validity of the test extract of P.D. in long term used.

The data showed in this model demonstrated that there was an decrease in blood sugar level in the extent of 28.06% & 33.17% in case of the test extract at the dose level of 100 mg/kg & 200mg/kg respectively on the 21<sup>st</sup> day of treatment.

While the standard drug on the same day bears 50.54% reduction at the same time tested dose levels & standard drug is P< 0.0001 on the 21<sup>st</sup> day.

The observed data results of the experiments suggested that the Ethanolic extract of P.D. able to maintain the hypoglycemic effect up to 21st day & no behavioural changes are observed during the treatment periods.

#### 6.7. Antidiabetic activity of ethanolic extract of leaves of PD in multidose treatment in alloxan induced Hyperglycemic rats in oral dose.

The purpose of the study is to confirm the antidiabetic effect of the test extract on longer duration of treatment.

In this model the test extract registered 52.44% & 57.58% of decrease of blood glucose level at the tested dose level of 100mg/kg & 200 mg/kg respectively on the 21<sup>st</sup> day of treatment. While the standard drug showed 60.83% reduction on the same day.

The test extract bears a significant P< 0.001 decrease value of blood sugar level starting from 7<sup>th</sup> day to 21<sup>st</sup> day at the both tested dose level while the standard drug showed the same rate significance which started from the 3<sup>rd</sup> day onwards. When compared with the solvent control.

The study further support the antidiabetic effect of the test extract whose effectiveness persist up to 21 days and the blood sugar level decreases gradually during the observed days, which presumed that the test extract, contains some antidiabetic active principle responsible for this effect.

The hypoglycemic active principle may act probably by initiating the release of insulin from the pancreatic cell of hyperglycemic animal (sulfonyl urea like effect)<sup>19</sup> the proper mechanism of action is further supported by OGTT in which glucose level reduces in response of test extract.

It is therefore, evident that hypoglycemic principle in the extract, exert a direct effect in diabetic rat probably by a mechanism similar to insulin. Which was substantiated by an extra pancreatic mode of action.

The experiment revealed that A.E. of P.D. (200mg/kg) significantly (P<0.001) decrease the glucose level on hyperglycemic animal. The glucose lowering activity observed in the diabetic animal may due to the stimulation of  $\beta$  cells in the pancreatic islets.

**Table 10:** Hypoglycemic activity of ethanolic extract of leaves of P.D in single dose treatment in glucose loaded hyperglycemic rats in oral route.(OGTT)

Group & treatment	Blood glucose levels(mg/dl)					%age decrease at the end of 8hrs
	-60min	0hrs	1hrs	2hrs	4hrs	
I. Solvent (saline Water)	104.16±2.21	107.83±.8	123.33±2.4	136±2.8	103.33±1.9	
II. Standard Glibenclamide(2.5mg/kg)	104.33±2.26	90±3.02***	78.83±2.2***	66±4.04***	55±4.3***	47.28
III. Test-1 (100mg/kg)	104.5±2.0	102.16±1.8	92.67±2.4***	89.16±2.08***	84.17±2.8***	19.45
IV. Test-2 (200mg/kg)	104±1.98	96.16±2.7**	88.84±2.7***	82.16±2.6***	79.5±2.2***	23.55
V. F Values		11.26 <sup>b</sup>	60.47 <sup>b</sup>	99.96 <sup>b</sup>	44.26 <sup>b</sup>	

Values are expressed in MEAN ± S.E.M of six animals. One Way ANOVA followed by Dunnet's t-test.

(F-value denotes statistical significance at (ap<0.05, bp<0.01)

(t-value denotes statistical significance at \*p<0.05, \*\*p<0.01 and \*\*\*p<0.001 respectively, in comparison to group-I.

**Table 11:** Hypoglycemic activity of ethanolic extract of leaves of P.D in multidose treatment in normoglycemic rats in oral dose

Group & treatment	Blood glucose levels(mg/dl)				%age decrease at the end of 8hrs
	0 <sup>th</sup> day	7 <sup>th</sup> day	14 <sup>th</sup> day	21 <sup>st</sup> day	
I. Solvent (saline Water)	104.66±2.85	104.83±2.07	104.66±1.7	105.5±2.2	
II. Standard Glibenclamide(2.5mg/kg)	106.16±2.28	86.16±3.13***	68.5±3.8***	52.5±2.6***	50.14
III. Test-1 (100mg/kg)	104.5±2.5	101.66±1.7	91.83±1.9**	75.17±4.5***	28.06
IV. Test-2 (200mg/kg)	104±2.0	98.83±1.8	85.34±2.6***	69.5±2.9***	33.17
V. F Values		13.01 <sup>b</sup>	31.62 <sup>b</sup>	47.2 <sup>b</sup>	

Values are expressed in MEAN ± S.E.M of six animals. One Way ANOVA followed by Dunnet's t-test.

(F-value denotes statistical significance at ap<0.05, bp<0.01)

(t-value denotes statistical significance at \*p<0.05, \*\*p<0.01 and \*\*\*p<0.001 respectively, in comparison to group-I.

**Table 12:** Antidiabetic activity of ethanolic extract of leaves of P.D in multidose treatment in alloxan induced Hyperglycemic rats in oral dose

Group & treatment	Blood glucose levels(mg/dl)					%age decrease at the end of 8hrs
	0 <sup>th</sup> day	3 <sup>rd</sup> day	7 <sup>th</sup> day	14 <sup>th</sup> day	21 <sup>st</sup> day	
I. Solvent (saline Water)	275.5±5.6	265±6.5	253.16±5.8	238.34±6.1	238.34±6.1	
II. Standard Glibenclamide(2.5mg/kg)	274.5±3.6	227.16±4.9***	182.5±6.7***	156.34±5.4***	107.5±4.07	60.83
III. Test-1 (100mg/kg)	279±4.5	263.83±4.6	233.16±5.7*	180.5±5.4***	132.67±4.02***	52.44
IV. Test-2 (200mg/kg)	280.16±4.8	263±4.3	226.33±6.5**	173.16±3.67***	118.83±2.9***	57.58
V. F Values		12.58 <sup>b</sup>	22.83 <sup>b</sup>	45.71 <sup>b</sup>	159.99 <sup>b</sup>	

Values are expressed in MEAN ± S.E.M of six animals. One Way ANOVA followed by Dunnet's t-test.

(F-value denotes statistical significance at (ap<0.05, bp<0.01)

(t-value denotes statistical significance at \*p<0.05, \*\*p<0.01 and \*\*\*p<0.001 respectively, in comparison to group-I.

### 6.8. Percentage loss in body weight in ethanolic extracts of leaves of PD in alloxan induced diabetic rats

The content of Table 13 indicates that there was decrease % loss in body wt. when measured on 3<sup>rd</sup>, 7<sup>th</sup>, 14<sup>th</sup> & 21<sup>st</sup> day of treatment in both tested dose levels.

The test extract at 100mg/kg possesses 6.72% loss of body wt. on 21<sup>st</sup> day. While on 3<sup>rd</sup> day the % loss of body wt. registered 12%. Similarly at dose level of 200mg/kg between, on 21<sup>st</sup> day measured 5.55% loss of body wt while on 3<sup>rd</sup> day % loss of body wt registered 10%.

However Glibenclamide (2.5mg/kg) reduces the body wt by 3.33% at the end of 21<sup>st</sup> days while the solvent control represented gradual increase in % loss of body wt. up to 21 days i.e. 21<sup>st</sup> day the loss was 25.33% where as on 3<sup>rd</sup> day the loss was 10%.

It has been reported that diabetes is a heterogeneous metabolic disorder in which one of the symptomatic character is loss in body wt. It has been stated that DM causes failure to use of glucose for energy leads to increased utilization & decreased storage of proteins responsible for reduction of body wt. essentially by depletion of the body proteins.<sup>17</sup>

The results of the study presented that the decreased loss of body wt. might be contributed by increased use of glucose by the tissue.

### 6.9. Effect of ethanolic extract of leaves of PD on the glycogen conc. in liver & kidney

The data presented in Table 14 demonstrated that the glycogen content of liver & kidney of DM animals treated with Ethanolic extract of leaves of PD & standard drug were significantly increased when compared with solvent control at the end of 21 days of treatment, even the test extract & standard drug approaches the content of glycogen towards normal glycogen level. It was reported that hyperglycemia results in decreased glucose utilization and glycogen synthesis.<sup>17</sup>

### 6.10. Effect of ethanolic extract of PD on lipid profile in Sub acute toxicity

The observation data should in Table 15 present significant (P<0.01) reduction of total cholesterol in serum in the groups treated with test extraction & standard drug when compared with solvent control.

The extent of reduction in both the close level of test extract was dose dependent. In case of triglycerides level & standard drug Glibenclamide (2.5mg/kg body wt.) reduces significantly (P<0.001) when compared with solvent control. All the above studies were carried out on the 21<sup>st</sup> day of treatment.

However, the value of HDL in case of test extract at tested dose levels & standard showed increased value than that of solvent control group with a significant level. While

the LDL & VLDL levels significantly (P<0.001) decrease in all drug treated groups when compared with solvent control.

It has been reported that in the untreated or under treated diabetic patients, hyperglyceridemia & hypercholesterolemia often occur due to the increase production of VLDL & due to unavailability of lipoprotein lipase which hydrolyses the triglycerides to VLDL because of insulin deficiency.<sup>20</sup>

As the test extract reduces VLDL, TC & TG. Hence it might be presumed that the test extract is responsible for the enhancement of the transcription of lipoprotein lipase, similar to insulin.

### 6.11. Effect of ethanolic extract of leaves of PD on hematological parameters in sub acute toxicity study

However, the diabetic rats treated with solvent showed a decrease value of RBC, WBC, & Hb. Content when compared with normal.

The hematological parameters exhibited in Table 16 showed that the animals treated with standard drug Glibenclamide & test extract in the tested dose levels bears normal value in RBC count & Hb. Count. Whereas the clotting time slightly elevated than normal. However, the diabetic rats treated with solvent showed a decrease value of RBC, WBC & Hb. Content when compared with normal.

The neutrophil count appears to nearly equal with that of normal value. The other hematological parameters like Eosinophil, lymphocyte & monocytes in case of the standard drug & test extract treated animals did not show any alteration.

Therefore, it might be suggested that the test extracts has no significant effect on the hematological parameters & is evident for the safely used at the ethanolic extract of P.D. leaves for a longer duration time.

### 6.12. Effect of ethanolic extract of leaves of PD on biochemical parameters in sub acute toxicity study

From the Table 17, the data showed ASAT & ALAT enzymes activity of alloxan induced diabetic rats showed significantly higher than normal rats. The test extract treated rats at tested dose levels had reduced enzymes activity when compared with diabetic rats.

The total protein & total albumin content serum was significantly lowered in diabetic rats when compared with solvent control but in the drug treated diabetic animals it returned to nearly normal.

The present study of experimental approach has been conducted for the test extract to elicit tonic response over an exposure period of 21 days. Alteration in the marker enzymes like ASAT, ALAT, ALP, liver & kidney glycogen content.

The total protein, lipid profile properties like TC, TG, HDL, LDL, VLDL of the extract were determined. The

**Table 13:** Percentage loss in body weight in ethanolic extract of leaves of P.D in alloxan induced diabetic rats

Group & treatment	Body weight loss in percentage			
	3 <sup>rd</sup> day	7 <sup>rd</sup> day	14 <sup>th</sup> day	21 <sup>st</sup> day
I. Solvent (saline Water)	10	16	18.66	25.33
II. Standard Glibenclamide(2.5mg/kg)	8	6.66	6.66	3.33
III. Test-1 (100mg/kg)	12	10.11	8.0	6.72
IV. Test-2 (200mg/kg)	10	8.23	6.38	5.55

**Table 14:** Effect of ethanolic extract of leaves of P.D on the glycogen conc. in liver & kidney

Group & treatment	Liver Glycogen mg/gm tissue	Kidney glycogen mg/gm tissue
I.Solvent (saline Water)	19.16±2.21	11.48±1.5
II.Standard Glibenclamide(2.5mg/kg)	33.66±2.76**	17.5±1.8*
III.Normal	35.67±2.7	19.4±1.4
IV. Test-I (100mg/kg)	28.46±1.19*	13.6±2.2
V. Test-II (200mg/kg)	31.8±4.4**	15.67±1.67
VI.F Values	5.21 <sup>b</sup>	3.14 <sup>a</sup>

Values are expressed in MEAN ± S.E.M of six animals. One Way ANOVA followed by Dunnet's t-test.

(F-value denotes statistical significance at (ap<0.05, bp<0.01)

(t-value denotes statistical significance at \*p<0.05, \*\*p<0.01 and \*\*\*p<0.001 respectively, in comparison to group-I.

**Table 15:** Effect of ethanolic extract of P.D on lipid profile in Sub acute toxicity

Group & treatment	Lipid Profile				
	TC mg/dl	TGmg/dl	HDLmg/dl	LDLmg/dl	VLDLmg/dl
I.Solvent (saline Water)	227.83±5.6	182±6.7	35.16±4.4	164.83±6.4	44.66±3.12
II.Standard Glibenclamide(2.5mg/kg)	161.34±4.9***	103.84±3.1***	47.67±3.2*	86.67±5.4***	23.83±2.8***
III.Normal	138.84±4.3	89.16±17	51.67±3.4	67.17±4.9	17.16±3.0
IV. Test-I (100mg/kg)	211.17±4.13	156.83±5.01**	39.17±4.8	146.67±5.8	32.67±2.6*
V. Test-II (200mg/kg)	203.47±5.7**	148.34±4.8***	41.17±3.7	136.83±4.4**	31.34±2.3**
VI.F Values	55.027 <sup>b</sup>	54.53 <sup>b</sup>	2.76 <sup>a</sup>	57.79 <sup>b</sup>	1351 <sup>b</sup>

Values are expressed in MEAN ± S.E.M of six animals. One Way ANOVA followed by Dunnet's t-test.

(F-value denotes statistical significance at (ap<0.05, bp<0.01)

(t-value denotes statistical significance at \*p<0.05, \*\*p<0.01 and \*\*\*p<0.001 respectively, in comparison to group-I.

**Table 16:** Effect of ethanolic extract of leaves of P.D on Hematological Parameters in sub acute toxicity study

Group & treatment	RBC (millions/ml)	WBC (millions/ml)	Hb (g/dl)	Clotting Time(min)	Neutrofil (%)	Esonophil (%)	Lymphocyte (%)	Monocyte (%)
I.Solvent (saline Water)	2.6±.42	7.8±.9	7.34±1.5	1.41±.009	21.16±3.7	4.75±.68	72.67±4.5	4.6±.7
II.Standard Glibenclamide(2.5mg/kg)	4.8±.6*	6.8±1.1	10.67±1.2	1.07±.08	27.83±2.3	1.91±.45***	69.84±3.3	2.8±.7
III.Normal	5.1±.6	7.1±1.24	11.84±1.4	1.05±.07	32.67±1.5	1.55±.07	68.67±3.3	1.9±.2
IV. Test-I (100mg/kg)	3.6±.8	6.1±.4	8.34±1.5	1.35±.07	24.34±2.1	2.6±.3**	71.34±4.1	3.67±.9
V. Test-II (200mg/kg)	4.16±.65	6.5±.7	8.84±1.2	1.25±.07	26.5±1.6	2.1±.4**	70.33±3.9	3.2±.3
VI.F Values	2.7 <sup>a</sup>	.45	1.6	3.87 <sup>a</sup>	3.1 <sup>a</sup>	8.4 <sup>b</sup>	0.15	2.17

Values are expressed in MEAN ± S.E.M of six animals. One Way ANOVA followed by Dunnet's t-test.

(F-value denotes statistical significance at (ap<0.05, bp<0.01)

(t-value denotes statistical significance at \*p<0.05, \*\*p<0.01 and \*\*\*p<0.001 respectively, in comparison to group-I.

**Table 17:** Effect of ethanolic extract of leaves of P.D on Biochemical parameters in sub acute toxicity study

Group & treatment	ASAT( $\mu$ l)	ALAT( $\mu$ l)	ALP( $\mu$ l)	Aibumin(gm/dl)	Total protein(gm.dl)
I.Solvent (saline Water)	38.16 $\pm$ 3.8	49.67 $\pm$ 4.09	291.5 $\pm$ 4.5	5.84 $\pm$ .79	4.75 $\pm$ .77
II.Standard Glibenclamide(2.5mg/kg)	25.17 $\pm$ 3.2*	31.34 $\pm$ 2.3**	270.84 $\pm$ 4.9*	4.2 $\pm$ .81	6.24 $\pm$ 1.03
III.Normal	22.84 $\pm$ 3.87	28.34 $\pm$ 3.4	262.17 $\pm$ 6.05	3.4 $\pm$ .37	8.08 $\pm$ .52
IV. Test-I (100mg/kg)	36.84 $\pm$ 3.87	39.84 $\pm$ 2.98*	280.17 $\pm$ 5.02	5.15 $\pm$ .79	5.14 $\pm$ .8
V. Test-II (200mg/kg)	30.67 $\pm$ 3.1	35.67 $\pm$ 2.24**	275.16 $\pm$ 4.49*	4.7 $\pm$ .67	5.67 $\pm$ .68
VI.F Values	3.86 <sup>a</sup>	7.24 <sup>b</sup>	4.67 <sup>b</sup>	1.71	2.78 <sup>a</sup>

Values are expressed in MEAN  $\pm$  S.E.M of six animals. One Way ANOVA followed by Dunnet's t-test.

(F-value denotes statistical significance at  $p < 0.05$ ,  $p < 0.01$ )

(t-value denotes statistical significance at \* $p < 0.05$ , \*\* $p < 0.01$  and \*\*\* $p < 0.001$  respectively, in comparison to group-I.

leaves of P.D. didn't produce any toxic or cells damaging effect to animals.

## 7. Conclusion

The ethanolic extract of leaves of *Phyllanthus distichus* results maximum yield value than that of petroleum ether extract, chloroform extract through successive extraction process. The ethanolic extract of leaves of *Phyllanthus distichus* showed maximum control in blood sugar in hyperglycemic wistar rats than other experimental extracts. The test extract also reduces the blood sugar level to a maximum extent in case of normal animals. Among the study of effects of AEPD leaves in both normoglycemic and hyperglycemic model through oral route. Toxicological study revealed that the AEPD leaves were safe and does not alter normal physiological and behavioral effect even at a higher dose level of 3000 mg/kg body weight.

Administration of AEPD significantly reduces the elevated glucose level in alloxan induced diabetic rats confirms its anti-diabetic activity. This also reduces normal glucose level, which reveals the hypoglycemic property. The whole protein, whole cholesterol, ASAT, ALAT, ALP enzyme activity of alloxan administered rats showed significantly higher than normal rats, the test extract treated rats significantly reduced the elevated levels. The results of the present investigation indicate that the AEPD leaves may have a place in the therapy of DM as anti-diabetic and/or hypoglycemic agent. The hypoglycemic and/or anti-diabetic effect of the test extract may be due to the influence on glycogenesis, glycogenolysis metabolic activity property of one or more of its constituents.

Thus it is concluded that the ethanolic extract of leaves of *Phyllanthus distichus* beneficial in lowering the blood sugar concentration and in management of other diabetic complications without any doubt.

## 8. Source of Funding

None.

## 9. Conflict of Interest

There is no conflict of interest among the authors.

## References

1. Al-Ahmed FAA, El-Denshary ESM, Zaki M, El-Sawaf HA, Abu-Jayyab AR. Interaction between diazepam and oral antidiabetic agents on serum glucose, insulin and chromium levels in rats. *Biosci Rep.* 1989;9:347–50.
2. Dash GK, Suresh P, Ganapati S. Studies on hypoglycaemic and wound healing activities of lantana camara linn. *J Nat Remedies.* 2001;1:105–10.
3. Ghosh M. Fundamentals of experimental pharmacology. 2nd ed. Calcutta: Scientific Book Agency; 1984.
4. Riley H, Spinks A. Biological assessment of tranquilisers. Part I. *J Pharmacol.* 1956;p. 657–71.
5. Jaiprakash R, Nagarani MA, Venkatraman BV. Effect of felodipine on cholinergic responses of the colonic smooth muscle of streptozotocin induced diabetic rats. *Indian J Exp Biol.* 1995;33(4):297–9.
6. Bergmeyer HU. Standardization of Enzyme Assays. *Clin Chem.* 1972;18(11):1305–11.
7. Richmond W. Preparation and properties of a cholesterol oxidase from *Nocardia* sp. and its application to the enzymatic assay of total cholesterol in serum. *Clin Chem.* 1972;19(12):1350–6.
8. Hendry R. Clinical chemistry. New York: Harper and Row Publishers; 1964.
9. Friedwald WT, Levy RI, Fredrickson DS. Estimation of the Concentration of Low-Density Lipoprotein Cholesterol in Plasma, Without Use of the Preparative Ultracentrifuge. *Clin Chem.* 1972;6(1):499–502.
10. Schettler G, Nussel E. Maßnahmen zur Prävention der Arteriosklerose. *Arb Med Soz Med Prav Med.* 1975;10:25.
11. Josephson G, Gyllensward C. The development of the protein fractions and of cholesterol concentration in the serum of normal infants and children. *Scand J Clin Lab Invest.* 1957;9(1):29–38.
12. Weichselbaum TE. An accurate and rapid method for the determination of proteins in small amounts of blood serum and plasma. *Am J Clin Pathol.* 1946;10:40–9.
13. Sood R. Medical laboratory technology, methods and interpretations. 5th ed.; 2003.
14. Morales MA, Jabbagy A, Terenzi HF. Mutations affecting accumulation of glycogen. *Fungal Genet Rep.* 1973;20. doi:10.4148/1941-4765.1830.
15. Ghai CL. A text book of practical physiology; 1998.
16. Kuikarni S. Handbook of experimental pharmacology. 2nd ed.; 1993.
17. Seth SD, Livingstone BI. 1999.
18. Chirdchupunseree H, Pramyothin P. Protective activity of phyllanthin in ethanol-treated primary culture of rat hepatocytes. *J Ethnopharmacol.* 2010;128(1):172–6.
19. Available from: [www.DiabetesIndia.com](http://www.DiabetesIndia.com).
20. Haardman JG, Limbird LE, Gilman AG. 1690.

**Author biography****Debashis Mohapatra** Assistant Professor**Sunita Mishra** Assistant Professor

**Cite this article:** Mohapatra D, Mishra S. Hypoglycemic and antidiabetic evaluation of leaves of phyllanthus distichus in normal and alloxan induced diabetic rats. *Indian J Pharm Pharmacol* 2020;7(3):189-202.