

Content available at: <https://www.ipinnovative.com/open-access-journals>

Indian Journal of Pharmacy and Pharmacology

Journal homepage: <https://www.ijpp.org.in/>

## Original Research Article

## To assess the rationality of pharmacotherapy of patient in clinical situations and validation of questionnaire tool used in rational pharmacotherapy

Avinash Vikram Turankar<sup>1,\*</sup>, Ashwita Uday Shetty<sup>1</sup>, Pramoda N<sup>1</sup>,  
Vijay Motiram Motghare<sup>1</sup>, Chetan Shankarrao Urade<sup>1</sup>

<sup>1</sup>Dept. of Pharmacology, Government Medical College, Nagpur, Maharashtra, India



## ARTICLE INFO

## Article history:

Received 07-11-2020

Accepted 25-11-2020

Available online 25-01-2021

## Keywords:

Rational pharmacotherapy

Rationality score

Questionnaire tool

## ABSTRACT

**Objectives:** 1. To provide a structured tool to assess rationality based on the 10 domains of Rationality score; 2. To find out the impact of the structured tool for judging the rationality score of Pharmacotherapy in various clinical situations e.g. High, Moderate and Low rationality score.

**Methodology:** A rationality scale was developed to check the coefficient of variation in each parameter of rationality, on a scale of 1 to 10, acronymed as R.A.T.I.O.N.A.L.I.T.Y.

Various clinical cases were discussed and treatment was criticised if any. These cases were then rated according to the rationality scale.

High Rationality - Score  $\geq 6$ , Average Rationality - Score  $> 4$  and  $< 6$  & Low Rationality - score  $\leq 4$ .

**Results:** For high rationality score, the top 5 parameter are availability of drugs, lab monitoring, rational combination, Protocol/Regime and if objectives of treatment fulfilled.

For average rationality scores, the top 5 parameters are Objective of treatment, Availability of drugs, Treatment correctness, lab monitoring and Protocol/Regime followed.

For low rationality, the lowermost 5 parameters were cost of therapy, irrational combination, lab monitoring, use of trade names instead of generic names and availability issues starting from the lowest.

**Conclusion:** Parameters like availability of the drugs, lab monitoring, presence or absence of irrational combinations are critical and common judgement parameters while judging if the therapy is rational or not.

© This is an open access article distributed under the terms of the Creative Commons Attribution License (<https://creativecommons.org/licenses/by/4.0/>) which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

### 1. Introduction

Pharmacotherapy, that is, drug therapy, plays an important role in people's health and their quality of life. However, there are increasing concerns with irrational drug use that can result in drug-related morbidity and mortality. These can increase the rate of adverse drug–drug interactions leading to additional cost and waste of valuable resources. The rapid increase in drug expenditure during the past decade, for example, has placed considerable pressure on healthcare systems.<sup>1</sup>

The concept of the rational use of medicines is an old one, dating as far back as 300 B.C, when the Greek

physician Herophilus said that “medicines are nothing in themselves, but are the very hands of god if employed with reason and prudence.”<sup>2,3</sup> According to World Health Organisation (WHO) definition, rational pharmacotherapy (RPT) requires that patients receive medications appropriate to their clinical needs, in doses that meet their own requirements, for an adequate period of time, and at the lowest cost to them and their community.<sup>3</sup> A multitude of factors have been found to lead to irrational use of drugs and these affect at various levels at which drugs are handled.<sup>4</sup> According to World Health Organisation (WHO) estimates, more than half of all medicines are prescribed, dispensed or sold inappropriately, and half of all patients fail to take medicines prescribed to them correctly.<sup>5,6</sup>

\* Corresponding author.

E-mail address: [aturankar@hotmail.com](mailto:aturankar@hotmail.com) (A. V. Turankar).

The word “ethics” is derived from the Greek word, *ethos*, which means custom or character. Ethics is an understanding of the nature of conflicts arising from moral imperatives and how best we may deal with them. It deals with the choices made and actions taken in relation to those choices.<sup>7,8</sup> It is concerned with the obligations of the doctors and the hospital to the patient along with other health professionals and society.<sup>9</sup> Medical ethics also deals with the choices made by society, the distribution of resources, and access to health care, and the dilemmas arising from them.<sup>7,10</sup>

Rationality, ethics are subjective issues, hence there is a wide variation and difficulty in the assessment of these parameters. There can be a positive and negative bias towards these parameters. There can be significant cognitive errors while assessing these aspects of research. But although these parameters are subjective and difficult to measure, they are meaningful and relevant in real life situations.

The current study is an effort to construct a tool for measuring the rationality pertaining to the pharmacotherapy of patient. A wide variety of literature regarding various aspects of rationality has been searched through and 10 important domains namely.

1. Regimen/Protocol – To check if cases follow standard guidelines or not
2. Availability of drugs – To check easy availability of drugs
3. Treatment correctness – To check whether drugs prescribed with actual dose, frequency, duration of therapy etc.
4. Information provided to patients or relatives – To check whether information has been given to patients/relatives regarding adverse effects of prescribed drugs, advice about therapy.
5. Objectives of management – To check if objectives of management are defined with respect to case or not
6. Need for drugs prescribed – To check for over prescription/under prescription
7. Amount/cost of therapy – To check if cost of regimen is up to the mark
8. Lab monitoring – To check if Lab monitoring is being done or not
9. Irrational combinations prescribed – To check rationality of prescribed treatment
10. Trade name/Generic names used – To check whether brand names used were identified as contents of rationality of pharmacotherapy of treatment. All these 10 domains can be summed up in acronym rationality.

## 2. Aims

To assess the rationality of pharmacotherapy of patient in clinical situations.

## 3. Objectives

### 3.1. Primary objective

To provide a structured tool to assess rationality based on the 10 domains of Rationality score mentioned above.

### 3.2. Secondary objectives

1. To find out the impact of the structured tool for judging the rationality score of Pharmacotherapy in various clinical situations e.g. High, Moderate and Low rationality score.
2. To assess the Validity of the questionnaire tool used in the assessment of the rational prescription behaviour of the clinicians.

## 4. Materials and Methods

A rationality scale was developed to check the coefficient of variation in each parameter of rationality. The parameters had been simplified on a scale of 1 to 10 based on subjective skills. In the middle of the scale 5 had been put as average. This was done because most of the subjective parameters were mentally judged on the basis of 0% to 100% and it was easy to quantify. These 10 parameters were acronymed as R.A.T.I.O.N.A.L.I.T.Y for the purpose of easy remembrance.

After formulating rationality scale, a trial version was tested. The post graduate students of Department of Pharmacology in a Tertiary Care Hospital presented various clinical cases as a part of their curriculum and criticised the treatment if any. These cases were discussed among professors, associate professors and lecturers and were then rated according to the rationality scale.

Scale of 0 to 4 were considered as low rationality, a scale of 4 to 6 were considered as average and scale of more than 6 were considered as good rationality. The domain scales were designed such that low score indicated low rationality and high score indicated high rationality judgement on that particular domain.

The analysis consisted of:

1. To find out the 5 most common parameters when the rating is judged as low and good
2. To find out the top 5 parameters in case of average rating i.e., between 4 to 6.

To assess the validity of the questionnaire tool the experts rated different aspects of the Questionnaire by answering 6 Questions rating 0 to 10 for each question in the online form. Faculty members with more than 5 years of experience from various colleges in the field of pharmacology have been considered to rate the scale.

## 5. Results

Overall we received 39 responses from the experienced pharmacologist who are the senior faculties of pharmacology. Approximately 90% of the faculties had an experience of more than 10 years.

**Table 1:** Score of individual questions

Question	Score (Mean $\pm$ SD)
1. Are all the elements or facets of the rationality or irrationality covered in the questionnaire?	8.718 $\pm$ 1.255
2. Are the questions in the questionnaire properly framed to understand each aspect of rationality?	8.513 $\pm$ 1.412
3. Is there clarity in framing of the questions?	8.282 $\pm$ 1.682
4. Is the rationality score in system properly designed?	8.590 $\pm$ 1.585
5. Can present rationality scoring system be used for predicting the rational prescription behaviour of physician?	8.667 $\pm$ 1.383
6. Do you feel that there is a correlation between the rational prescription behaviour and present scales?	7.897 $\pm$ 2.150

**Table 2:** Experience distribution

Work Experience	Number of Participants
5 to 10 years	5 (12.8%)
11 to 20 years	11 (28.2%)
20 to 30 years	19 (48.7%)
More than 30 years	4 (10.3%)

**Table 3:** Gender distribution

Total participants	Male	Female
39	23 (59%)	16 (41%)

## 6. Discussion

The overall rationality score was analysed and the data was divided into three categories of less than and equal to 4, between 4 and 6 and greater than and equal to 6.

For higher rationality score, that is score greater than and equal to 6 which means when the assessors judged the rationality score to be high, the top five parameter are availability of the drugs, lab monitoring, presence of rational combination, if the pharmacotherapy was as per Regime or protocol and if the therapy had an objective of treatment.

For average rationality scores, the top 5 parameters are if the therapy had objective of treatment, the availability of drugs, treatment correctness, lab monitoring and last if the pharmacotherapy was as per resume of protocol.

For low rationality score, that is less than and equal to 4 the lowermost 5 parameters were, cost of therapy, presence of irrational combination in the therapy, if lab monitoring was done properly, use of trade names in place of generic names and availability issues starting from the lowest.

We can conclude from the above results that some parameters are given critical importance while judging the rationality score as high, low or average. These common parameters are availability of the drugs, lab monitoring, presence or absence of irrational combinations.

The other important judgement criteria were whether the treatment protocol and regimens were followed or not, presence of objectives for the treatment, treatment correctness and use of trade names while prescribing.

Regarding the validity of the scale, the senior faculties were of the opinion that all the elements and facets of the rationality or irrationalities are covered in the questionnaire to the extent of 87%. The framing of the questions were proper to the extent of 85%, the clarity of the questions was to the extent of 82%, the design of the questions in the rationality was proper to the extent of 85%.


The senior faculty were of the opinion that the same system can be used for predicting rational prescription behaviour of the physician to the extent of 87%. There was 79% feeling that there is a correlation between the rational prescription behaviour and the present irrationality scale.

On the whole on majority of the parameters the scale was valid to the extent of 80 to 87% on various different parameters.


Reliability of rationality score over time and within assessors needs to be evaluated in future studies.

Despite this encouraging judgement and response by the senior faculties they came up with various promising suggestions.


One of the suggestion related with the question number 4 was that information of the prescription should be given to the patient and their relatives in their vernacular language. There was one very good suggestion by a senior faculty that the clinicians should also be involved in this prescription rationality analysis. So that it becomes realistic and pragmatic. One recommendation was that regular rationality checks like this will help the cause of rational pharmacotherapy in the long run. Similarly it was suggested that this particular scale is good for detecting the gross rational behaviour than one of incidences which can be exceptions. One of the senior faculty suggested to weigh these various different parameters differently because rationality cannot be scored equally. Some of the important parameters may have different weightage than the others, so it should be taken into account while formulating such a scale.



**Fig. 1:** High rationality score - score greater than and equal to 6 (\*Parameters arranged in descending order of importance. \*Top 5 parameters to be considered)



**Fig. 2:** Average rationality score – score greater than 4 and less than 6 (\*Parameters arranged in descending order of importance \*Top 5 parameters to be considered)



**Fig. 3:** Low rationality score – less than and equal to 4 (\*Parameters arranged in descending order of importance \*Lowest 5 parameters to be considered)

## 7. Conclusion

Parameters like availability of the drugs, lab monitoring, presence or absence of irrational combinations are critical and common judgement parameters while judging the therapy as rational or not. In future these rationality score parameters may be useful for judging the rationality of the prescription behaviour of physician or the hospital etc., but it needs to be evolved over a period of time and scientifically validated as well.

In the absence of any standardised validated rational pharmacotherapy assessment scale, the present scale cannot be correlated with any gold standard scale. The above findings pertaining to validity are standalone with respect to present scale.

The researchers are keen to incorporate these various suggestions into their subsequent versions of the rational pharmacotherapy scales in the times to come.

## 8. Source of Funding

None.

## 9. Conflict of Interest

None.

## References

1. World Health Organization. The world medicines situation report 2011. The world medicines situation report 2011.
2. Asenso RO, Agyeman AA. Irrational Use of Medicines—A Summary of Key Concepts. *Pharm (Basel)* . 2016;4:35. doi:10.3390/pharmacy4040035.
3. Ambwani S, Mathur AK. Chapter-2 Rational Drug Use. *Health Administrator*. 2006;19(1):5–7.

4. Shivhare SC, Kunjwani HK, Manikrao AM, Bondre AV. Drugs Hazards and Rational Use of Drugs: A. *J Chem Pharm Res*. 2010;2(1):106–12.
5. Lukali V, Michelo C. Factors associated with irrational drug use at a district hospital in Zambia: patient record-based observations. *Med J Zambia*. 2015;42(1):25–30.
6. Laing R, Ruredzo R. The essential drugs programme in Zimbabwe: new approaches to training. *Health Policy Plan*. 1989;4:229–34. doi:10.1093/heapol/4.3.229.
7. Avasthi A, Ghosh A, Sarkar S, Grover S. Ethics in medical research: General principles with special reference to psychiatry research. *Indian J Psychiatry*. 2013;55(1):86. doi:10.4103/0019-5545.105525.
8. Thatte U. Basic Principles of Clinical Research and Methodology. *Ethical issues in Clinical Research*. 2007;p. 58–73.
9. Markose A, Krishnan R, Ramesh M. Medical ethics. *J Pharm Bioallied Sci*. 2016;8(1):1–4.
10. World Health Organization. Operational guidelines for ethics committees that review biomedical research. World Health Organization; 2000.

## Author biography

**Avinash Vikram Turankar**, Associate Professor

**Ashwita Uday Shetty**, Junior Resident

**Pramoda N**, Junior Resident

**Vijay Motiram Motghare**, Professor and Head

**Chetan Shankarrao Urade**, Ex-Assistant Professor

**Cite this article:** Turankar AV, Shetty AU, Pramoda N, Motghare VM, Urade CS. To assess the rationality of pharmacotherapy of patient in clinical situations and validation of questionnaire tool used in rational pharmacotherapy. *Indian J Pharm Pharmacol* 2020;7(4):251-255.