

Content available at: <https://www.ipinnovative.com/open-access-journals>

Indian Journal of Pharmacy and Pharmacology

Journal homepage: <https://www.ijpp.org.in/>

## Original Research Article

## Evaluation of potential antidiabetic activity of *abelmoschus esculentus* and evidence of possible pharmacokinetic interaction with metformin

Debalina Sardar<sup>1</sup>, Raja Chakraverty<sup>2,\*</sup>, Pranabesh Chakraborty<sup>3</sup><sup>1</sup>Dept. of Pharmacology, Bengal School of Technology (A College of Pharmacy), West Bengal, India<sup>2</sup>Dept. of Pharmacology, Institute of Postgraduate Medical Education and Research, Kolkata, India<sup>3</sup>Dept. of Pharmaceutical and Medical Science and Technology, Maulana Abul Kalam Azad University of Technology, West Bengal, India

## ARTICLE INFO

## Article history:

Received 22-11-2021

Accepted 09-12-2021

Available online 05-01-2022

## Keywords:

Abelmoschus esculentus

Metformin

Antagonism

Diabetes mellitus

## ABSTRACT

The study is aimed at the evaluation of potential activity of *Abelmoschus esculentus* and possible interaction with metformin in animal Models of Diabetes Mellitus. Study objectives include study the anti-diabetic effect of *Abelmoschus esculentus* for Diabetes Mellitus in animal models and also to study the effect of *Abelmoschus esculentus* with metformin and explore any interaction. Plant material was collected (*Abelmoschus esculentus*) followed by extraction of plant materials (*Abelmoschus esculentus*) Exudate collection of *Abelmoschus esculentus* and activity test study was done (acute toxicity study, according to standard OECD guidelines) Experimental animals were divided into groups. Dosing was done for 28 days. Biochemical parameters were studied. Histopathology studies are done. Results showed that in this study administrations of *Abelmoschus esculentus* extract (2000mg/kg body weight) Metformin with *Abelmoschus esculentus* extract (5mg/kg b.w. and 2000mg/kg body weight and Metformin 5mg/kg body weight decreased elevated blood glucose levels significantly from first to fourth week compared to diabetic control rats and showed minimal safety concerns.

This is an Open Access (OA) journal, and articles are distributed under the terms of the [Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License](https://creativecommons.org/licenses/by-nc-sa/4.0/), which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

For reprints contact: [reprint@ipinnovative.com](mailto:reprint@ipinnovative.com)

## 1. Introduction

Okra (*Abelmoschus esculentus*), also known as “lady’s fingers,” is a green flowering plant. Okra belongs to the same plant family as hibiscus and cotton.<sup>1–6</sup> The term “okra” most commonly refers to the edible seedpods of the plant. Okra has long been favored as a food for the health conscious people. It contains nutrients like potassium, vitamin B, vitamin C, folic acid, and calcium. It’s low in calories and has high dietary fiber content. Recently, a new benefit of including okra in your diet is being considered. Okra has been suggested to help manage blood sugar in cases of type 1, type 2,

and gestational diabetes.

The study is aimed at the Evaluation of Potential Activity of *Abelmoschus esculentus* and Possible Interaction with Metformin in Animal Models of Diabetes Mellitus. Study objectives include study the anti-diabetic effect of *Abelmoschus esculentus* for Diabetes Mellitus in animal models and also to study the effect of *Abelmoschus esculentus* with metformin and explore any interaction.

## 2. Materials and Methods

Plant material was collected (*Abelmoschus esculentus*) followed by extraction of plant materials (*Abelmoschus esculentus*) Exudate collection of *Abelmoschus esculentus* Pre- Processing Activity test study was done (acute toxicity study, according to standard OECD guidelines). Animals

\* Corresponding author.

E-mail address: [rchakraborty20@yahoo.com](mailto:rchakraborty20@yahoo.com) (R. Chakraverty).

are divided into five groups. The experimental animals received drugs for 28 days.<sup>1,7-11</sup> Biochemical studies are checked. Histopathology studies are done. Results, data were collected and estimated Results and data are estimated and graphs are plotted.

### 2.1. Raw plant material collection

500 gm of fresh pod or fruits of *Abelmoschus esculentus* were collected from the local market of Sugandhya, Delhi road, Chinsurah, West Bengal and a sample was authenticated by the Botanical survey of India, Shibpur as *Abelmoschus esculentus* which was archived for future reference. (BSI/Plant spec./2016).<sup>12-15</sup>

### 2.2. Chemicals

Metformin HCL, Glucose, Sucrose, streptozotocin, nicotinamide, DPPH, Accu-check Glucometer, Auto-analyser (Rovonic) Experiments were performed on a total of 15 Albino Wistar rats weighing about 100-200 gm of either sex. They were maintained in colony cages (3 rats per cage) at an ambient temperature of 25-27°C with 12 hours light and dark cycles having proper ventilation in the room. They were fed normal diets purchased commercially from the vendor and water. The rats were allowed to acclimatize to the laboratory environment for one week and were randomly divided into groups for experiments.<sup>16,17</sup>

Preparation of viscous water-soluble fraction of *Abelmoschus esculentus*: 500 gm of fresh pods or fruits of okra (*Abelmoschus esculentus*) were collected. Then, the pods were thoroughly washed with distilled water, cut into small slices by a sharp knife. The sliced pods were immersed into distilled water in a beaker. The mixture was then stirred gently for 10-15 minutes with a glass rod. After that, it was filtered using a thin layer of cotton to remove the insoluble matters and filtrate was collected. The amount of filtrate was measured.

### 2.3. Induction of diabetes

Diabetes was induced in rats by a single intra peritoneal injection of streptozotocin (45 mg/kg body weight) dissolved into citrate buffer. 15 minutes later of streptozotocin injection, nicotinamide injection (110 mg/kg body weight) was induced by the same route. 21 days later, plasma glucose concentration was determined in blood samples obtained from rats. Rats with diabetic (>200 mmol/L) were included in the study and the rats which no diabetic (<100mmol/L) were excluded from the experiment.

### 2.4. Experimental design

Rats were divided into five groups. Each group contained three Wistar albino rats (n = 3 per groups). The groups are

divided as Normal control, Diabetic control (diabetic but were not under the treatment, DC) Diabetic with Metformin only; Metformin control (MC) Diabetic with WSF of *Abelmoschus esculentus* (WFC) Diabetic with Metformin and WSF of *Abelmoschus esculentus* (DMF)

### 2.5. Acute toxicity study

Acute toxicity study of *Abelmoschus esculentus* extract was determined as per Organization for Economic Cooperation and Development guidelines (OECD) 423.

Table 1:

| Concentration of DPPH+ ascorbic acid | Absorbance of test group | Mean value |
|--------------------------------------|--------------------------|------------|
| 20                                   | 0.573                    |            |
| 40                                   | 0.583                    |            |
| 60                                   | 0.585                    | 0.588      |
| 80                                   | 0.590                    |            |
| 100                                  | 0.610                    |            |




Fig. 1: Blood sugar levels in experimental groups




Fig. 2: Liver function tests (ALT and AST)




Fig. 3: Antidiabetic experimental groups




Fig. 4: Serum albumin

Five (05) mice were used for the acute toxicity study. For acute toxicity study mice were treated with the 2000mg/kg body weight extract of okra by oral route. After the oral administration of okra extract, animals were observed individually at least once during the first 30 minutes, periodically during the first 24 hours, with special attention given during first 4 hours, and 14 days regularly observed for toxicity determination of extract of *Abelmoschus esculentus*.

Dosing protocol using 5mg/kg body weight of the standard drug metformin was weighted and dissolved in sterile distilled water. It was given to the Metformin Control (MC) group. 0.4ml/L animal body weight extract of Okra (*Abelmoschus esculentus*) was measured and it was given to

the Diabetic with WSF of *Abelmoschus esculentus* (WFC) group. *Abelmoschus esculentus* extract was given to the Diabetic rats with metformin and WSF of *Abelmoschus esculentus* (DMF) group. The Diabetic Control group and the normal Control group were not treated by the test drugs. The groups were fed with normal diet and water was given ad libitum.

### 3. Statistical Analysis

All the data were expressed as mean + SEM were evaluated by one-way analysis of variance (ANOVA) followed by Drunnett's test for multiple comparisons using prism Graphpad version 5.0 and values of  $P < 0.05$  were considered as statistically significant.

### 4. Results

Results of the In-Vitro Anti-Oxidant activity of Ascorbic acid by DPPH free radical scavenging method: Calculation of Antioxidant activity test group is summarised in Table 1.

### 5. Discussion

There were no significant changes in the mice body weight. The dose was 2000 mg/kg. The oral administration of okra (*Abelmoschus esculentus*) extract did not show any toxicity signs and mortality up to 14 days during acute toxicity study. The extract was found to be safe at the dose level of 2000mg/kg body weight. There was no toxic effect, no lethal effect on animals in all groups after 14 days of observation. LD50 higher than 15 gm/kg b.w. is categorized "Practically Non-Toxic". In this study, administrations of *Abelmoschus esculentus* (okra) extract (2000mg/kg b.w.), Metformin with *Abelmoschus esculentus* extract (5mg/kg b.w. and 2000mg/kg b.w.) and Metformin (5mg/kg b.w.) decreased elevated blood glucose levels significantly from first to fourth week compared to diabetic control rats.

### 6. Conclusion

In this study administrations of *Abelmoschus esculentus* (okra) extract (2000 mg/kg body weight led to reduction of fasting and post prandial glucose levels, Metformin with *Abelmoschus esculentus* extract (5mg/kg body weight. And 2000 mg/kg body weight and metformin 5mg/kg body weight decreased elevated blood glucose levels significantly from first to fourth week compared to diabetic control rats and showed minimal safety concerns.

### 7. Source of Funding

None.

### 8. Conflict of Interest

None.

## References

1. Khatun H, Rahman A, Biswas M, Islam AU. Watersoluble Fraction of *Abelmoschus esculentus* L Interacts with Glucose and Metformin Hydrochloride and Alters Their Absorption Kinetics after Co administration in Rats. *ISRN Pharm*. 2011;2011:1–5. doi:10.5402/2011/260537.
2. Centers for Disease Control and Prevention. Centers for Disease Control and Prevention. Diabetes Report Card 2014. Atlanta, GA: Centers for Disease Control and Prevention, US Dept. of Health and Human Services; 2015. ; 2014.
3. World Health Organisation. Global status report on NCDs; 2010. Available from: <https://www.who.int/news-room/fact-sheets/detail/noncommunicable-diseases>.
4. Krentz AJ, Bailey CJ. Oral antidiabetic agents: current role in type 2 diabetes mellitus. *Drugs*. 2005;65(3):385–411.
5. Pelikanova T. Treatment of diabetes in metabolic syndrome. *Vnitř Lek*. 2009;55(7-8):637–45.
6. Moise MM, Benjamin LM, Doris TM, Dalida KN, Augustin NO. Role of Mediterranean diet, tropical vegetables rich in antioxidants, and sunlight exposure in blindness, cataract and glaucoma among African type 2 diabetics. *Int J Ophthalmol*. 2012;5(2):231–7. doi:10.3980/j.issn.2222-3959.2012.02.23.
7. Gul MZ, Bhakshu LM, Ahmad F, Kondapi AK, Qureshi IA, Ghazi IA. Evaluation of *Abelmoschus moschatus* extracts for antioxidant, free radical scavenging, antimicrobial and antiproliferative activities using in vitro assays. *BMC Complement Altern Med*. 2011;11(1):64. doi:10.1186/1472-6882-11-64.
8. Knekt P, Kumpulainen J, Jarvinen R, Rissanen H, Heliövaara M, Reunanen A. Flavonoid intake and risk of chronic diseases. *Am J Clin Nutr*. 2002;76(3):560–8. doi:10.1093/ajcn/76.3.560.
9. Fayaz SM, Kumar S, Rajanikant VS. Finding Needles in a Haystack: Application of Network Analysis and Target Enrichment Studies for the Identification of Potential Anti-Diabetic. *PLoS ONE*. 2014;9(11):112. doi:10.1371/journal.pone.0112911.
10. Sabitha V, Ramachandran S, Naveen KR, Panneerselvam K. Antidiabetic and antihyperlipidemic potential of *Abelmoschus esculentus* (L.) Moench. In streptozotocin-induced diabetic rats. *J Pharm Bioallied Sci*. 2011;3(3):397–402. doi:10.4103/0975-7406.84447.
11. Fan S, Zhang Y, Sun Q, Yu L, Li M, Zheng B. Extract of okra lowers blood glucose and serum lipids in high-fat diet-induced obese C57BL/6 mice. *J Nutr Biochem*. 2014;25(7):702–9. doi:10.1016/j.jnutbio.2014.02.010.
12. Tian ZH, Miao FT, Zhang X, Wang QH, Lei N, Guo LC. Therapeutic effect of okra extract on gestational diabetes mellitus rats induced by streptozotocin. *Asian Pac J Trop Med*. 2015;8(12):1038–42. doi:10.1016/j.apjtm.2015.11.002Get.
13. Choi HN, Kang MJ, Lee SJ, Kim JI. Ameliorative effect of myricetin on insulin resistance in mice fed a high-fat, high-sucrose diet. *Nutr Res Pract*. 2014;8(5):544–9. doi:10.4162/nrp.2014.8.5.544.
14. Ong KC, Khoo HE. Effects of myricetin on glycemia and glycogen metabolism in diabetic rats. *Life Sci*. 2000;25(14):1695–705. doi:10.1016/s0024-3205(00)00758-x.
15. Ozcan F, Ozmen A, Akkaya B, Aliciguzel Y, Aslan M. Beneficial effect of myricetin on renal functions in streptozotocin-induced diabetic diabetes. *Clin Exp Med*. 2012;12(4):265–72. doi:10.1007/s10238-011-0167-0.
16. Kandasamy N, Ashokkumar N. Protective effect of bioflavonoid myricetin enhances carbohydrate metabolic enzymes and insulin signaling molecules in streptozotocin-induced diabetic nephrotoxic rats. *Toxicol Appl Pharmacol*. 2014;279(2):173–85. doi:10.1016/j.taap.2014.05.014.
17. Song Y, Manson JE, Buring JE, Sesso HD, Liu S. Associations of dietary flavonoids with risk of type 2 diabetes, and markers of insulin resistance and systemic inflammation in women: a prospective study and cross-sectional analysis. *J Am Coll Nutr*. 2005;24(5):376–84. doi:10.1080/07315724.2005.10719488.

## Author biography

**Debalina Sardar**, Assistant Professor

**Raja Chakraverty**, Assistant Professor

**Pranabesh Chakraborty**, Director

**Cite this article:** Sardar D, Chakraverty R, Chakraborty P. Evaluation of potential antidiabetic activity of *abelmoschus esculentus* and evidence of possible pharmacokinetic interaction with metformin. *Indian J Pharm Pharmacol* 2021;8(4):263-266.